

*Systematic Review*

# Efficacy and Safety of Rechallenge with BRAF/MEK Inhibitors in Advanced Melanoma Patients: A Systematic Review and Meta-Analysis

Jonathan N. Priantti <sup>1</sup> , Maysa Vilbert <sup>2,3</sup> , Thiago Madeira <sup>4</sup>, Francisco Cezar A. Moraes <sup>5</sup>, Erica C. Koch Hein <sup>2,3,6</sup>, Anwaar Saeed <sup>7</sup> and Ludimila Cavalcante <sup>8,\*</sup> 

- <sup>1</sup> School of Medicine, Federal University of Amazonas—UFAM, Manaus 69020-160, AM, Brazil
- <sup>2</sup> Princess Margaret Cancer Centre, University Health Network, Toronto, ON M5G 2M9, Canada
- <sup>3</sup> Division of Medical Oncology, Department of Medicine, University of Toronto, Toronto, ON M5S 1A8, Canada
- <sup>4</sup> School of Medicine, Federal University of Minas Gerais—UFMG, Belo Horizonte 30130-100, MG, Brazil
- <sup>5</sup> School of Medicine, Federal University of Pará—UFPA, Belém 66075-110, PA, Brazil
- <sup>6</sup> Department of Hematology and Oncology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago 8331150, Chile
- <sup>7</sup> Department of Medicine, Division of Hematology and Oncology, UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA 15213, USA
- <sup>8</sup> Department of Medical Oncology, Novant Health Cancer Institute, Charlotte, NC 28204, USA
- \* Correspondence: ludi.cavalcante@gmail.com



**Citation:** Priantti, J.N.; Vilbert, M.; Madeira, T.; Moraes, F.C.A.; Hein, E.C.K.; Saeed, A.; Cavalcante, L. Efficacy and Safety of Rechallenge with BRAF/MEK Inhibitors in Advanced Melanoma Patients: A Systematic Review and Meta-Analysis. *Cancers* **2023**, *15*, 3754. <https://doi.org/10.3390/cancers15153754>

Academic Editors: Rachid Skouta, Armando Varela-Ramirez, Elisa Robles-Escajeda, Blanca E. Ruiz-Medina and Patricia Talamás-Rohana

Received: 25 June 2023

Revised: 20 July 2023

Accepted: 23 July 2023

Published: 25 July 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Simple Summary:** Approximately 50% of patients with melanoma harbor a BRAF mutation and are eligible for targeted therapy with BRAF/MEK inhibitors (BRAFi/MEKi). Despite a response rate of nearly 70%, more than half of the patients will experience disease progression within a year due to tumor resistance. Rechallenging patients with BRAFi/MEKi has emerged as an alternative for improving response and survival outcomes. This systematic review and meta-analysis aims to investigate the efficacy and safety of this strategy in patients with advanced melanoma.

**Abstract:** This systematic review and meta-analysis aims to evaluate the efficacy and safety of rechallenging advanced melanoma patients with BRAFi/MEKi. Seven studies, accounting for 400 patients, were included. Most patients received immunotherapy before the rechallenge, and 79% underwent rechallenge with the combination of BRAFi/MEKi. We found a median progression-free survival of 5 months and overall survival of 9.8 months. The one-year survival rate was 42.63%. Regarding response, ORR was 34% and DCR 65%. There were no new or unexpected safety concerns. Rechallenge with BRAFi/MEKi can improve outcomes in advanced melanoma patients with refractory disease. These findings have significant implications for clinical practice, particularly in the setting of progressive disease in later lines and limited treatment options.

**Keywords:** advanced melanoma; targeted therapy; rechallenge; BRAF/MEK inhibitors; MAPK inhibitors

## 1. Introduction

Melanoma represents a significant health problem and economic burden [1,2]. According to recent data from the National Center for Health Statistics, 97,610 new cases of melanoma will be diagnosed in the US in 2023, with approximately 8000 deaths [3]. The incidence has been overall stable, with the rate decreasing in men and increasing in older women by 1% per year [3,4]. The highest incidence is seen in Australia/New Zealand, followed by Western Europe, North America, and Northern Europe [2]. Worldwide, the mortality rate is disproportionately higher in transitioning countries, likely due to difficulties in accessing the newest medications [2,5]. The insufficient evidence for screening as a secondary prevention for the early detection of melanoma patients contributes to the disease

burden [6,7]. In addition, some patients experience disease progression or are diagnosed with advanced disease, leading to poorer outcomes [8,9].

Approximately a decade ago, the prognosis of metastatic melanoma patients was dismal, and half would die within a year of diagnosis [10]. Available treatments at that time were ineffective, with low and short-lived response with chemotherapy or significant toxicity from agents, such as high-dose interleukin 2 (HD IL-2) [11–15]. Fortunately, the discovery of driver mutations and immune evasion mechanisms led to the development of targeted therapies and immune checkpoint inhibitors [16], and their approval by the Food and Drug Administration (FDA) and other regulatory agencies was considerable in this regard [11,17]. Several targeted therapies were developed, showing superiority compared to the chemotherapy dacarbazine, which was considered a reasonable approach before the FDA approval of Vemurafenib (BRAF inhibitor) and Ipilimumab (anti-CTLA-4) in 2011 [18,19]. The advances in treatments led to a large increase of 258% in the number of individuals living with metastatic melanoma in the last two decades [20].

The BRAF gene encodes different RAF protein isoforms that activate serine/threonine protein kinases in the mitogen-activated protein kinase (MAPK) signaling pathway [21,22]. BRAF mutations are oncogenic drivers that stimulate cell proliferation and growth through this pathway, augmenting the capacity of tumors to progress and spread [23–25]. Roughly 50% of melanomas harbor a BRAF mutation, especially the variant V600E, suitable for targeted therapy with BRAF inhibitors [22,26].

Vemurafenib was the first drug used in this context, and since its discovery, other BRAF inhibitors, such as Dabrafenib and Encorafenib, have become available [27–29]. Concurrently, the MAP kinase (MAPK) pathway was meticulously explored, and an important resistance mechanism to BRAFi, the downstream activation of MEK, was identified [11,24]. MEK inhibitors (MEKi) were then incorporated to the treatment landscape. Moreover, clinical studies have demonstrated that the combination of BRAF and MEK inhibition enhances the MAPK pathway blockade, resulting in better and consistent outcomes than BRAFi alone [30–39]. Surprisingly, this combination also exhibits a better safety profile than either BRAFi or MEKi monotherapy [39–41], also showing improvement in health-related quality of life in unresectable III/IV stage melanoma patients [42]. However, despite the advantages offered by targeted therapy with BRAFi/MEKi combination, more than 50% of patients still progress within a year of treatment due to the reactivation of the MAPK pathway through alternative pathways [11,18,43].

Immunotherapy with immune checkpoint inhibitors has been the treatment of choice when targeted therapy fails due to its long-term survival benefit, especially in complete responders [18,44–46]. Unfortunately, almost half of the patients will not respond due to primary resistance mechanisms, and most responders may ultimately progress [18,47]. Further treatment options for these patients include a clinical trial, when available, or chemotherapy [48]. More recently, alternatives, such as treatment beyond progression and rechallenge, have stood out [49–52]. The latter has shown promising results in melanoma, consisting of re-exposure to BRAFi/MEKi after a period on another treatment or a drug holiday [26]. The BRAFi/MEKi-free interval, in theory, allows for sensitive tumor cells to grow, and a subsequent rechallenge could hypothetically induce responses again and improve survival outcomes [53,54]. Given the emerging clinical evidence for this practice, our systematic review and meta-analysis aims to investigate the efficacy and safety of rechallenging advanced BRAF-mutant melanoma patients with MAPK inhibitors.

## 2. Materials and Methods

### 2.1. Eligibility Criteria

Inclusion in this meta-analysis was restricted to studies that met all the following eligibility criteria: (1) randomized and non-randomized clinical trials and prospective and retrospective observational cohort studies; (2) including advanced melanoma patients who previously progressed on BRAFi  $\pm$  MEKi in 1st or 2nd line therapy; and (3) patients rechallenged with BRAFi/MEKi after other systemic treatments or a drug holiday. In addition,

studies were only included if they reported any of the clinical outcomes of interest: survival outcomes, response to BRAFi/MEKi rechallenge, and safety. We excluded studies with the following: (1) no outcomes of interest; (2) case reports, series of cases, or case-control studies; (3) no BRAFi/MEKi rechallenge; or (4) studies with overlapping populations.

## 2.2. Search Strategy and Data Extraction

We systematically searched PubMed, Scopus, and Cochrane Central Register of Controlled Trials databases, and the congresses websites of the American Society of Clinical Oncology (ASCO) publications, European Society for Medical Oncology (ESMO), and American Association for Cancer Research (AACR) from inception to November 2022. The following search terms were used: “melanoma”, “rechallenge”, “disease progression”, “progressed disease”, “re-challenge”, “BRAF/MEK”, “BRAF”, “MEK”, “trametinib”, “cobimetinib”, “binimetinib”, “vemurafenib”, “dabrafenib”, and “encorafenib”. The references from all included studies and previous systematic reviews were also searched manually for any additional studies. Three authors (J.N.P., T.M., and F.C.A.M.) independently extracted the data following the predefined search criteria and quality assessment. If there was no unanimity about including any study, a fourth author decided its eligibility (M.V.).

## 2.3. Endpoints and Subanalysis

Efficacy outcomes included objective response rate (ORR), disease control rate (DCR), progression-free survival (PFS), overall survival (OS), and one-year overall survival rate (1-year OS). ORR included patients with partial or complete response, whereas DCR included patients with stable disease, partial or complete response. Response was evaluated by investigator–clinical assessment or by RECIST (Response Evaluation Criteria in Solid Tumors) criteria version 1.1, according to each study’s own criteria. In the safety analysis, adverse events were characterized based on the Common Terminology Criteria for Adverse Events (CTCAEs) version 4.0. The prespecified analyses of prognostic factors included subgroups of patients with elevated lactate dehydrogenase (LDH) and metastasis to the central nervous system. We sought all results compatible with our outcomes of interest in each study. Then, we built a table containing the study characteristics, interventions, and results and compared them against our planned outcomes to define the groups for each synthesis.

## 2.4. Quality Assessment

Non-randomized interventional studies were assessed through the Risk Of Bias In Non-randomized Studies of Interventions (ROBINS-I) tool [55], which contains seven domains and categorizes studies as having low, moderate, serious, critical, or unclear risk of bias. The quality assessment of observational studies was performed using the Newcastle–Ottawa Scale (NOS), in which studies are scored on a 0 to 9 scale according to selection, comparability, and exposure criteria [56].

## 2.5. Statistical Analysis


The systematic review and meta-analysis were performed in accordance with the Cochrane Collaboration and the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement guidelines (PRISMA Checklist, Supplementary Tables S1 and S2) [57]. We conducted a proportional meta-analysis pooling the data with the function “metaprop” and “pool.median”, included in the packages “meta”, “metafor”, and “metamedian” in R for efficacy outcomes [58]. We did not report OS and PFS in hazard ratios (HRs) due to the absence of this information in most studies. Thus, we pooled the medians of these variables through the method proposed by McGrath et al. [59]. The HRs with 95% confidence interval (CI) from multivariate analyses were used to assess prognostic factors, such as elevated LDH and the presence of brain metastasis, using the function “metagen”, also included in the “meta” package in R. Cochran’s Q test and  $I^2$  statistics were used to evaluate heterogeneity;  $p$ -values inferior to 0.10 and  $I^2 > 25\%$  were considered

significant for heterogeneity. A DerSimonian and Laird random-effects model was applied to all analyses. We used RStudio (Posit Software, PBC, version 2022.12.0 + 353) for statistical analysis. The study protocol was registered in the International Prospective Register of Systematic Reviews database—PROSPERO (registration number: CRD42022375952).

### 3. Results

#### 3.1. Characteristics of the Included Studies and Patients

The initial literature search generated 1858 results, as detailed in Figure 1. After removing duplicate records and ineligible studies, 34 remained and were thoroughly reviewed based on inclusion criteria. Of these, a total of seven studies were included: one phase II clinical trial and six observational cohort studies comprising 400 patients [60–66]. A full list of the excluded studies can be found in the Supplementary Materials, Table S3.



**Figure 1.** PRISMA flow diagram of the study screening and selection.


The population was characterized by BRAF-mutant advanced melanoma patients, who had targeted therapy in the first or second line of treatment, with either the combination of BRAFi/MEKi or BRAFi alone. Most patients (83%, 333/400) underwent an interval treatment with immunotherapy before being rechallenged with targeted therapy, and 10% (40/400) went on a drug holiday. During the rechallenge, 79% patients (317/400) were treated with the combination of BRAFi/MEKi, whereas 21% (83/400) received BRAFi alone. The baseline characteristics of the included studies are shown in Table 1, and the summary of the treatments administered during the first targeted therapy, interval treatment, and rechallenge are presented in Figure 2. The description of the best response during the first targeted therapy (TT) exposure (in the first/second line of treatment) and the best response

with TT rechallenge in each study are presented in the Supplementary Materials, Figure S1 and Table S4.

**Table 1.** Baseline characteristics of the population in the included studies.

Study ID	N	Age <sup>†</sup> , y	TT in 1st/2nd Line with BRAFi + MEKi (%)	IT with ICI (%)	Interval between TT and RC <sup>†</sup> (Months)	RC with BRAFi + MEKi (%)	ECOG (0–1, ≥2) (%)	LDH (Normal, ≥ULN) (%)	CNS Disease at RC (yes, no) (%)
Atkinson (2020) [60]	90	61	80	100	NA	93	60, 28	30, 51	49, 46
Cybulska-Stopa (2020) [61]	51	56	68	100	8.6	96	78, 22	22, 76	59, 41
Persa (2021) [62]	48	57	79	75	4	83	NA	40, 60	50, 50
Roux (2015) [63]	10	52.4	0	80	NA	10	50, 50	70, 30	60, 40
Schreuer (2017) [64]	25	54.7	64	100	6.1	100	80, 20	72, 28	68, 32
Tietze (2018) [65]	60	56	32	67	3.4	68	63, 32	43, 57	60, 40
Valpione (2017) [66]	116	51.9	35	71	7.7	66	61, 22	39, 55	44, 56

N: number. y: year. TT: targeted therapy. IT: interval treatment. ICI: immune checkpoint inhibitors. RC: rechallenge. BRAFi: BRAF inhibitor. MEKi: MEK inhibitor. LDH: lactate dehydrogenase. ULN: upper limit of normality. CNS: central nervous system. NA: not available. <sup>†</sup> Median.




**Figure 2.** Summary of the treatments administered during the first targeted therapy (TT) exposure (A), during the interval period (B), and the TT rechallenge period (C). During the first TT, most patients (58%) received the combination of BRAFi plus MEKi, and 42% received BRAFi alone. A total of 83% of patients received immunotherapy during the interval treatment, whereas 10% were on a drug holiday. In the rechallenge setting, almost 80% of patients received the combination of BRAFi plus MEKi, while 21% were treated with BRAFi alone.

### 3.2. Efficacy and Safety of Rechallenging Advanced Melanoma Patients with Targeted Therapy

In a pooled analysis including all seven studies, the ORR was 34.25% (95% CI 28.5 to 40.0) with no significant heterogeneity seen among the studies ( $I^2 = 25\%$ ,  $p = 0.24$ ), as shown in Figure 3A. The prevalence of DCR, presented in Figure 3B, was 65.01% (95% CI 57.31 to 72.72) with a high level of inter-study heterogeneity ( $I^2 = 59\%$ ,  $p = 0.02$ ).

Exploring the prevalence of DCR further in a sensitivity analysis, we found that studies with a median BRAFi/MEKi-free interval of six months or more achieved a significantly higher DCR rate than those with a shorter BRAFi/MEKi-free interval (<6 months). The DCR rate was 68% versus 56%, respectively ( $p < 0.01$ ), with a low heterogeneity in each subgroup ( $I^2$  of 0% and 12%, respectively), as illustrated in Figure 4.




**Figure 3.** Forest plots showed an ORR of 34.25% (A), a DCR of 65.01% (B), and a 1-year OS rate of 42.63% (C) [60–66].

The median progression-free survival (mPFS) during rechallenge with BRAFi/MEKi was 5 months (95% CI 4 to 5.9), based on a pool of medians from five studies. The median OS was 9.8 months (95% CI 9.3 to 20.4), according to data from four studies. The proportion of patients alive in 1 year was 42.63% (95% CI 30.25 to 55.02), with considerable variability among studies ( $I^2 = 70\%$ ,  $p = 0.01$ ), as depicted in Figure 3C.

We conducted pooled analyses of hazard ratios from three studies assessing PFS and two studies evaluating OS, comparing (1) the presence or absence of brain metastasis and (2) normal versus elevated LDH levels. However, only some studies provided univariate and multivariate analyses evaluating prognostic factors associated with survival outcomes. Consequently, the findings of our combined sub-analyses were also limited. The analyses did not yield statistically significant results. The forest plots can be found in the Supplementary Materials, Figure S2.

Regarding safety, there were no new or unexpected adverse events. Two studies reported that the safety profile during TT rechallenge was very similar to the first TT treatment [60,63], while in one study [66], patients reported a better tolerance during rechallenge. There were no deaths related to treatment. The most frequent toxicities were fever and rash.





**Figure 4.** Subgroup analysis of DCR during targeted therapy (TT) rechallenges according to the median BRAFi/MEKi-free interval. Patients with a BRAFi/MEKi-free interval  $\geq 6$  months had a DCR of 68.3%, significantly higher than patients with a BRAFi/MEKi-free interval  $< 6$  months, with a DCR of 56.2% ( $p < 0.01$ ) [60–66].

### 3.3. Quality Assessment

Our results consist of information collected from one clinical trial and six observational studies. The non-randomized clinical trial was considered to have a moderate risk of bias. All other studies scored 7 out of 9 in NOS, except for one observational study scoring 6 out of 9 due to lack of information on confounding factors [64]. Individual appraisal of non-randomized interventional studies and observational studies can be seen, respectively, in Supplementary Figure S3 and Supplementary Table S5.

In the visual inspection of the two funnel plots from the outcomes of DCR and 1-year OS rate, we found a predominantly symmetrical pattern of the number of studies on each side and their disposition around the central axis, as shown in Figure S4. However, two studies were located slightly outside the plot in each analysis, which may represent differences in patient selection and intervention. Aiming to explore it further, we ran Egger's regression test, which showed no evidence suggesting significant publication bias in either of the two outcomes (Supplementary Figure S4A,B; DCR:  $z = 1.7615$ ,  $p = 0.0782$ ; and 1-year OS rate:  $z = -0.4770$ ,  $p = 0.6355$ ).

We performed leave-one-out sensitivity analyses by systematically removing each study from the pooled estimates in two variables with high heterogeneity: the prevalence of DCR and the prevalence of 1-year OS rate (Supplementary Figure S5A,B, respectively). Applying the leave-one-out test did not substantially alter the results. Additionally, when analyzing the influence of some studies on DCR, Roux et al. [34], a small-sized study of nine patients, had the most significant benefit in DCR, although it was the study with the worst 1-year OS rate. When excluded from some of our analyses, we found this study contributed significantly to the high heterogeneity ( $I^2 = 0$  when removed).

## 4. Discussion

In this systematic review and meta-analysis of seven studies, including 400 patients, we assessed the efficacy and safety of rechallenging BRAF-mutant advanced melanoma

patients with BRAFi/MEKi. The main findings related to this therapeutic strategy include the following: (1) BRAFi/MEKi rechallenge is a promising palliative treatment, with more than two-thirds of patients achieving disease control mainly due to partial response and stable disease; (2) an interval therapy longer than 6 months was significantly associated with a greater DCR; (3) objective response was seen in more than one-third of patients; (4) median PFS was 5 months; (5) median OS was 9.8 months, and 43% of patients were alive at 1 year; (6) the presence of brain metastasis or elevated LDH did not show a statistically significant association with an increased risk of progression or death; and (7) patients had a good tolerance, without unexpected adverse events during TT rechallenge.

Since the BRIM-3 trial with Vemurafenib [27,67] and the BREAK-3 trial with Dabrafenib [28,68], BRAF inhibitors have shown greater benefit in improving outcomes for BRAF-mutant unresectable or metastatic melanoma patients compared to chemotherapy with dacarbazine. Moreover, the combination of BRAF and MEK inhibitors has yielded even more promising results, as demonstrated in the COMBI-D phase III trial, combining Dabrafenib with Trametinib [32–34,69]. This combination enhanced outcomes by improving the efficacy and reducing the toxicity compared to BRAF monotherapy [70]. Additional clinical trials have also substantiated this approach's efficacy, as exemplified by the coBRIM trial (Vemurafenib plus Cobimetinib) and the 5-year update of the COLUMBUS trial (Encorafenib plus Binimetinib), which showed consistent efficacy and safety of the BRAFi/MEKi combination [30,31,35–37,71]. Unfortunately, most patients will eventually progress despite the encouraging initial responses to treatment [34,43]. This acquired resistance leads to a gradual loss of clinical response during treatment, primarily due to the reactivation of the MAPK/ERK downstream pathway or at the level of the BRAF mutation itself [70,72–75].

Moreover, recent studies have shown other potential mechanisms involved in response and resistance, such as tumor cell differentiation, epigenetic reprogramming, the production of bioregulatory factors as self-protection against host responses and therapies, and the neuro-endocrine and immune system modulation by melanin pigmentation and melanogenesis to promote tumor growth [76–79]. In the genetic realm, the upregulation of mutated genes, such as MITF and the receptor tyrosine kinase AXL, as well as downregulation of the tumor suppressor gene PTEN, are associated with BRAF inhibitor resistance [80–82]. Additionally, phenotype switching, which confers heterogeneity in MAPK pathway dependence, contributes to sensitivity or resistance to TT and is an area of opportunity for future drug development [83–85].

In all studies included in our research, TT was the first-line treatment for metastatic melanoma. Although BRAF inhibition can enhance antigen expression, facilitates T-cell cytotoxicity, and creates a favorable tumor microenvironment, potentially augmenting the efficacy of immunotherapy [86–88], recent investigations revealed that resistance to BRAF/MEK inhibitors induces an immune-evasive tumor microenvironment [89,90]. This cross-resistance to immunotherapy in melanoma cells, was characterized by the appearance of non-functional CD103+ dendritic cells in the tumor microenvironment, as a result of MAPK pathway reactivation [90]. As a result, impaired antigen presentation hinders effective T-cell responses and potentially decreases the efficacy of sequential immunotherapy [90]. In contrast, immunotherapy may potentially enhance the response to targeted therapy in BRAF-mutated melanoma, thereby prolonging tumor regression durability [91].

The most current clinical data support the use of ICI as the preferred first-line treatment for BRAF-mutated melanoma [92]. Studies, such as SECOMBIT and DREAMseq, which aimed to determine the optimal initial treatment for these patients, demonstrated that immunotherapy following progression on first-line BRAF/MEK inhibitors yields lower response rates and worse survival compared to the reverse treatment sequence [93–95]. All these data support the interaction between two types of treatments with completely different mechanisms of action enhancing or decreasing each other's activity. About 83% of patients included in the meta-analysis were on immunotherapy during the interval treatment, potentially creating a favorable setting for TT rechallenge or being influenced by the immune-evasive state promoted after first-line TT. We could not properly assess



the influences of targeted therapy on immunotherapy response nor the influences of immunotherapy on TT rechallenge response due to insufficient data, thereby directing this issue for further investigation.

While immunotherapy has been established as the first-line option for treatment-naïve metastatic melanoma patients, including those harboring a BRAF mutation [92–94], patients who progress on a targeted therapy and later on immunotherapy, or the inverse order, still have limited treatment options and poor prognosis [88,96–100]. Given this landscape, several case series emerged demonstrating encouraging activity with the re-introduction of targeted therapy (TT) in patients with prior progression on BRAFi/MEKi who went on drug holiday or interval treatment [49–52]. Therefore, rechallenging patients with targeted therapy could overcome the resistance mechanisms related to the previous progression to a BRAFi alone or the combination of BRAFi/MEKi [49–54].

In our analysis, prolonged interval therapy from the first TT exposure until rechallenge (greater than 6 months) was associated with a DCR of almost 70%, which is significantly higher than in patients with a short interval therapy (<6 months, DCR of 56%). This difference may be explained by a reversible mechanism of resistance. Tumors are heterogeneous, and the prevalence of distinct intra-tumoral cellular clones may change as the cell replicates and acquires new mutations, or according to the variations in the tumor microenvironment, or by external pressures, such as systemic or local therapies [75,101–103]. Hence, after exposure and initial response, some tumoral clones may develop resistance to TT, allowing them to proliferate and survive, leading to disease progression. Once TT is discontinued and the patient switches to a different interval therapy or drug holiday, external pressure changes, allowing other tumoral clones to arise and proliferate. This process may result in a reversion of BRAFi/MEKi resistance due to the growth of sensible tumor clones, resulting in a de novo response to TT [103].

A network meta-analysis of 15 RCTs evaluating systemic therapy for previously untreated advanced BRAF-mutated melanoma demonstrated that patients treated with BRAF and MEK inhibitors achieved higher ORR compared with BRAF alone (odds ratio of 2.00), and both had superior objective responses compared to other treatments [104]. In line with that, combined TT is the standard of care in current clinical practice due to what is known about resistance to single-agent BRAF inhibition. TT combination is administered to very symptomatic patients with a high burden of disease in the first-line therapy, aiming for a faster response than immunotherapy [11,92]. In contrast, some of the included studies in our meta-analysis did not use combination TT as a first-line treatment, leading to some of the limitations described below.

Of note, in our analysis, about 42% of patients received BRAFi alone and not combination BRAFi/MEKi as the initial TT exposure prior to rechallenge, which may have positively influenced the responses to therapy found in our study. In these patients, it is possible that MEK sensitivity remained intact through a lack of external pressure and those patients could potentially benefit more when treated with the combination of BRAFi/MEKi in the rechallenge setting. Conversely, 21% of patients were also treated with BRAFi alone in the rechallenge setting. Due to lack of patient-level data, we could not explore this hypothesis further or evaluate the magnitude of influence this had on our results. Currently, combined TT is the standard of care for BRAF-mutant melanoma patients [105,106], and the response rates with TT rechallenge in clinical practice may be somewhat lower than what was reported in this paper.

Our population consisted of 53% (164/310) of patients with brain metastasis. We analyzed the presence of brain metastasis as a prognostic factor in addition to the limited data available for this subgroup and did not find a significant association with OS and PFS. The brain is a frequent site of recurrence in BRAF-mutant melanoma patients across different ethnicities [107–109]. Studies have indicated a higher incidence of brain metastases in patients treated with BRAF/MEK inhibitors compared to those treated with immunotherapy [107,110]. This highlights the importance of continuous surveillance and management strategies for brain metastases [107,110]. One interesting question is if the

recurrence patterns after rechallenge are similar to the first TT exposure. Specifically, understanding brain metastasis response and recurrence would inform us about the central nervous system penetration of TT rechallenge and the need of further local therapy. The information regarding progressive sites of disease after rechallenge was not available in our studies. Thus, we were unable to analyze whether there is a difference in the incidence of CNS disease recurrence and progression between first-line TT therapy and rechallenge.

The LDH level is a well-known factor associated with prognosis in melanoma patients [111–113]. In our population, 48.5% (194/400) of patients had high LDH levels. Yet, we could not demonstrate a significant association of LDH with OS and PFS due to a lack of survival data in this subgroup among the studies. Interestingly, regardless of these high-risk features (elevated LDH and brain metastasis), our systematic review and meta-analysis demonstrated benefits in tumor response with TT rechallenge for all population.

None of the studies in our meta-analysis reported any unexpected adverse events when rechallenging advanced melanoma patients with BRAFi/MEKi. This underscores the well-known safety profile of this treatment strategy, which was also observed in the first-line therapy and can be applied to the rechallenge setting [39,114].

## 5. Limitations

Our study has some limitations. We did not pool data on time-to-event survival outcomes, such as PFS and OS, due to the constraints of missing information in the included studies. We managed this by reporting PFS and OS as the median of medians. Even though most of our studies were observational cohort studies and may have selection bias influencing their results, we were able to demonstrate tumor response to TT rechallenge with low heterogeneity in our primary or sensitivity analyses. Notwithstanding, our study presented new and relevant data concerning the benefit of adding a palliative treatment with TT rechallenge for BRAF-mutant metastatic melanoma patients.

## 6. Conclusions

To the best of our knowledge, this is the first systematic review and meta-analysis to pool available data on rechallenging BRAF-mutant advanced melanoma patients with MAPK inhibitors. Our findings indicate that most of patients can benefit from TT rechallenge by achieving the partial control or stability of their disease. Better outcomes were observed in patients with a TT-free interval greater than six months. Additionally, the median overall survival was almost ten months in the third-line setting in all population analyses, including patients with brain metastasis and high LDH.

Our study provided important evidence that rechallenge with BRAFi/MEKi should be considered as a palliative treatment option for patients who progress on MAPKi and had an interval period with or without other treatments. Further research will be important to investigate the patterns of recurrence and progression during rechallenge and the best sequencing strategy to optimize patient response and survival.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/cancers15153754/s1>. Table S1. PRISMA 2020 Checklist. Table S2. PRISMA 2020 for abstract checklist. Table S3. Full list of the excluded studies after a comprehensive analysis. Table S4. Description of the best response during the first targeted therapy (TT) exposure (in the 1st/2nd line of treatment) and the best response with TT rechallenge in each study. Table S5. Individual study appraisal for assessing the quality of non-randomized studies in the meta-analyses using the Newcastle–Ottawa Scale (NOS). Figure S1. Summary of the best overall responses (%) during targeted therapy (TT) 1st exposure and rechallenge. Figure S2. Prognostic factors associated with OS (A,B) and PFS (C,D). Figure S3. Bias assessment through ROBINS-I for non-randomized interventional studies. Figure S4. Funnel plots for publication bias analyses of DCR (A) and 1-year OS rate (B). Figure S5. Leave-one-out test analyses of DCR (A) and 1-year OS rate (B).

**Author Contributions:** Conceptualization, J.N.P. and M.V.; methodology, J.N.P. and M.V.; provision of study materials, J.N.P., T.M., F.C.A.M. and M.V.; collection and congregation of data, J.N.P., T.M., F.C.A.M. and M.V.; data analysis and interpretation, J.N.P. and M.V.; writing—original draft preparation, J.N.P., M.V., E.C.K.H., L.C., A.S., F.C.A.M. and T.M.; writing—review and editing, J.N.P., M.V., E.C.K.H., L.C., A.S., F.C.A.M. and T.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** The data for this study were systematically collected and organized into a comprehensive database. Access to the data can be granted upon request from the corresponding author.

**Conflicts of Interest:** The authors J.N.P., T.M., F.C.A.M. and M.V., declare no conflict of interest. E.C.K.H.: Consulting or Advisory Role (Novartis). Speaker’s Bureau (Novartis and MSD). Travel, Accommodation, and Expenses (Pfizer, Novartis, and Roche Pharma AG). Research Funding (Novartis, funding paid to Dr. Koch Hein Institution for support of a melanoma registry in Chile). L.C.: Employment (Actuate Therapeutics). Consulting or Advisory Role (Pliant Therapeutics, Janssen, and CDR-Life). Stock and Other Ownership Interests (Actuate Therapeutics). A.S. reports research grants (to institution) from AstraZeneca, Bristol Myers Squibb, Merck, Clovis, Exelixis, Actuate Therapeutics, Incyte Corporation, Daiichi Sankyo, Five prime therapeutics, Amgen, Innovent biologics, Dragonfly therapeutics, KAHN medical, and Biontech, and advisory board fees from AstraZeneca, Bristol Myers Squibb, Exelixis, Pfizer, and Daiichi Sankyo.

## Abbreviations

BRAFi: BRAF inhibitor. DCR: Disease control rate. FDA: Food and Drug Administration. IL-2: Interleukin-2. MAPK: MAP kinase. MEKi: MEK inhibitor. ORR: Objective response rate. OS: Overall survival. PFS: Progression-free survival. TT: Targeted therapy.

## References

1. Parra, L.M.; Webster, R.M. The Malignant Melanoma Market. *Nat. Rev. Drug Discov.* **2022**, *21*, 489–490. [\[CrossRef\]](#)
2. Arnold, M.; Singh, D.; Laversanne, M.; Vignat, J.; Vaccarella, S.; Meheus, F.; Cust, A.E.; de Vries, E.; Whiteman, D.C.; Bray, F. Global Burden of Cutaneous Melanoma in 2020 and Projections to 2040. *JAMA Dermatol.* **2022**, *158*, 495. [\[CrossRef\]](#)
3. Siegel, R.L.; Miller, K.D.; Wagle, N.S.; Jemal, A. Cancer Statistics, 2023. *CA Cancer J. Clin.* **2023**, *73*, 17–48. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Cronin, K.A.; Scott, S.; Firth, A.U.; Sung, H.; Henley, S.J.; Sherman, R.L.; Siegel, R.L.; Anderson, R.N.; Kohler, B.A.; Benard, V.B.; et al. Annual Report to the Nation on the Status of Cancer, Part 1: National Cancer Statistics. *Cancer* **2022**, *128*, 4251–4284. [\[CrossRef\]](#) [\[PubMed\]](#)
5. Fundytus, A.; Sengar, M.; Lombe, D.; Hopman, W.; Jalink, M.; Gyawali, B.; Trapani, D.; Roitberg, F.; De Vries, E.G.E.; Moja, L.; et al. Access to Cancer Medicines Deemed Essential by Oncologists in 82 Countries: An International, Cross-Sectional Survey. *Lancet Oncol.* **2021**, *22*, 1367–1377. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Mangione, C.M.; Barry, M.J.; Nicholson, W.K.; Chelmsow, D.; Coker, T.R.; Davis, E.M.; Donahue, K.E.; Jaén, C.R.; Kubik, M.; Li, L.; et al. Screening for Skin Cancer. *JAMA* **2023**, *329*, 1290. [\[CrossRef\]](#)
7. de Vere Hunt, I.; Lester, J.; Linos, E. Insufficient Evidence for Screening Reinforces Need for Primary Prevention of Skin Cancer. *JAMA Intern. Med.* **2023**, *183*, 509. [\[CrossRef\]](#) [\[PubMed\]](#)
8. Melanoma Survival Rates | Melanoma Survival Statistics. Available online: <https://www.cancer.org/cancer/melanoma-skin-cancer/detection-diagnosis-staging/survival-rates-for-melanoma-skin-cancer-by-stage.html> (accessed on 23 February 2023).
9. Sadeq, M.A.; Ashry, M.H.; Ghorab, R.M.F.; Afify, A.Y. Causes of Death among Patients with Cutaneous Melanoma: A US Population-Based Study. *Sci. Rep.* **2023**, *13*, 10257. [\[CrossRef\]](#) [\[PubMed\]](#)
10. Berk-Krauss, J.; Stein, J.A.; Weber, J.; Polsky, D.; Geller, A.C. New Systematic Therapies and Trends in Cutaneous Melanoma Deaths Among US Whites, 1986–2016. *Am. J. Public Health* **2020**, *110*, 731–733. [\[CrossRef\]](#)
11. Curti, B.D.; Faries, M.B. Recent Advances in the Treatment of Melanoma. *N. Engl. J. Med.* **2021**, *384*, 2229–2240. [\[CrossRef\]](#)
12. Hersh, E.M.; Del Vecchio, M.; Brown, M.P.; Kefford, R.; Loquai, C.; Testori, A.; Bhatia, S.; Gutzmer, R.; Conry, R.; Haydon, A.; et al. A Randomized, Controlled Phase III Trial of Nab-Paclitaxel versus Dacarbazine in Chemotherapy-Naïve Patients with Metastatic Melanoma. *Ann. Oncol.* **2015**, *26*, 2267–2274. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Falkson, C.I.; Ibrahim, J.; Kirkwood, J.M.; Coates, A.S.; Atkins, M.B.; Blum, R.H. Phase III Trial of Dacarbazine versus Dacarbazine with Interferon Alpha-2b versus Dacarbazine with Tamoxifen versus Dacarbazine with Interferon Alpha-2b and Tamoxifen in

- Patients with Metastatic Malignant Melanoma: An Eastern Cooperative Oncology Group Study. *J. Clin. Oncol.* **1998**, *16*, 1743–1751. [[CrossRef](#)] [[PubMed](#)]
14. Middleton, M.R.; Grob, J.J.; Aaronson, N.; Fierlbeck, G.; Tilgen, W.; Seiter, S.; Gore, M.; Aamdal, S.; Cebon, J.; Coates, A.; et al. Randomized Phase III Study of Temozolomide versus Dacarbazine in the Treatment of Patients with Advanced Metastatic Malignant Melanoma. *J. Clin. Oncol.* **2000**, *18*, 158. [[CrossRef](#)] [[PubMed](#)]
  15. Hauschild, A.; Garbe, C.; Stolz, W.; Ellwanger, U.; Seiter, S.; Dummer, R.; Ugurel, S.; Sebastian, G.; Nashan, D.; Linse, R.; et al. Dacarbazine and Interferon  $\alpha$  with or without Interleukin 2 in Metastatic Melanoma: A Randomized Phase III Multicentre Trial of the Dermatologic Cooperative Oncology Group (DeCOG). *Br. J. Cancer* **2001**, *84*, 1036–1042. [[CrossRef](#)] [[PubMed](#)]
  16. Pasquali, S.; Kefford, R.; Chiarion Sileni, V.; Nitti, D.; Rossi, C.R.; Pilati, P.; Mocellin, S. Systemic Treatments for Metastatic Cutaneous Melanoma. *Cochrane Database Syst. Rev.* **2014**, *2*, CD011123. [[CrossRef](#)]
  17. Scott, E.C.; Baines, A.C.; Gong, Y.; Moore, R.; Pamuk, G.E.; Saber, H.; Subedee, A.; Thompson, M.D.; Xiao, W.; Pazdur, R.; et al. Trends in the Approval of Cancer Therapies by the FDA in the Twenty-First Century. *Nat. Rev. Drug Discov.* **2023**, 1–16. [[CrossRef](#)]
  18. Thompson, J.F.; Kefford, R.; Stevens, G.; Scolyer, R. *Melanoma*; Riker, A.I., Ed.; Springer International Publishing: Cham, Switzerland, 2018; ISBN 978-3-319-78309-3.
  19. Lopes, J.; Rodrigues, C.M.P.; Gaspar, M.M.; Reis, C.P. Melanoma Management: From Epidemiology to Treatment and Latest Advances. *Cancers* **2022**, *14*, 4652. [[CrossRef](#)]
  20. Gallicchio, L.; Devasia, T.P.; Tonorezos, E.; Mollica, M.A.; Mariotto, A. Estimation of the Number of Individuals Living with Metastatic Cancer in the United States. *JNCI J. Natl. Cancer Inst.* **2022**, *114*, 1476–1483. [[CrossRef](#)]
  21. Davies, H.; Bignell, G.R.; Cox, C.; Stephens, P.; Edkins, S.; Clegg, S.; Teague, J.; Woffendin, H.; Garnett, M.J.; Bottomley, W.; et al. Mutations of the BRAF Gene in Human Cancer. *Nature* **2002**, *417*, 949–954. [[CrossRef](#)]
  22. Dankner, M.; Rose, A.A.N.; Rajkumar, S.; Siegel, P.M.; Watson, I.R. Classifying BRAF Alterations in Cancer: New Rational Therapeutic Strategies for Actionable Mutations. *Oncogene* **2018**, *37*, 3183–3199. [[CrossRef](#)]
  23. Alqathama, A. BRAF in Malignant Melanoma Progression and Metastasis: Potentials and Challenges. *Am. J. Cancer Res.* **2020**, *10*, 1103–1114.
  24. Proietti, I.; Skroza, N.; Bernardini, N.; Tolino, E.; Balduzzi, V.; Marchesiello, A.; Michelini, S.; Volpe, S.; Mambrin, A.; Mangino, G.; et al. Mechanisms of Acquired BRAF Inhibitor Resistance in Melanoma: A Systematic Review. *Cancers* **2020**, *12*, 2801. [[CrossRef](#)] [[PubMed](#)]
  25. Akbani, R.; Akdemir, K.C.; Aksoy, B.A.; Albert, M.; Ally, A.; Amin, S.B.; Arachchi, H.; Arora, A.; Auman, J.T.; Ayala, B.; et al. Genomic Classification of Cutaneous Melanoma. *Cell* **2015**, *161*, 1681–1696. [[CrossRef](#)] [[PubMed](#)]
  26. Serra, F.; Barruscotti, S.; Dominioni, T.; Zuccarini, A.; Pedrazzoli, P.; Chiellino, S. Treatment Following Progression in Metastatic Melanoma: The State of the Art from Scientific Literature to Clinical Need. *Curr. Oncol. Rep.* **2021**, *23*, 84. [[CrossRef](#)]
  27. Chapman, P.B.; Hauschild, A.; Robert, C.; Haanen, J.B.; Ascierto, P.; Larkin, J.; Dummer, R.; Garbe, C.; Testori, A.; Maio, M.; et al. Improved Survival with Vemurafenib in Melanoma with BRAF V600E Mutation. *N. Engl. J. Med.* **2011**, *364*, 2507–2516. [[CrossRef](#)]
  28. Hauschild, A.; Grob, J.-J.; Demidov, L.V.; Jouary, T.; Gutzmer, R.; Millward, M.; Rutkowski, P.; Blank, C.U.; Miller, W.H.; Kaempgen, E.; et al. Dabrafenib in BRAF-Mutated Metastatic Melanoma: A Multicentre, Open-Label, Phase 3 Randomised Controlled Trial. *Lancet* **2012**, *380*, 358–365. [[CrossRef](#)]
  29. Delord, J.-P.; Robert, C.; Nyakas, M.; McArthur, G.A.; Kudchakar, R.; Mahipal, A.; Yamada, Y.; Sullivan, R.; Arance, A.; Kefford, R.F.; et al. Phase I Dose-Escalation and -Expansion Study of the BRAF Inhibitor Encorafenib (LGX818) in Metastatic BRAF-Mutant Melanoma. *Clin. Cancer Res.* **2017**, *23*, 5339–5348. [[CrossRef](#)]
  30. Larkin, J.; Ascierto, P.A.; Dréno, B.; Atkinson, V.; Liskay, G.; Maio, M.; Mandalà, M.; Demidov, L.; Stroyakovskiy, D.; Thomas, L.; et al. Combined Vemurafenib and Cobimetinib in BRAF-Mutated Melanoma. *N. Engl. J. Med.* **2014**, *371*, 1867–1876. [[CrossRef](#)]
  31. Ascierto, P.A.; McArthur, G.A.; Dréno, B.; Atkinson, V.; Liskay, G.; Di Giacomo, A.M.; Mandalà, M.; Demidov, L.; Stroyakovskiy, D.; Thomas, L.; et al. Cobimetinib Combined with Vemurafenib in Advanced BRAFV600-Mutant Melanoma (CoBRIM): Updated Efficacy Results from a Randomised, Double-Blind, Phase 3 Trial. *Lancet Oncol.* **2016**, *17*, 1248–1260. [[CrossRef](#)]
  32. Long, G.V.; Stroyakovskiy, D.; Gogas, H.; Levchenko, E.; de Braud, F.; Larkin, J.; Garbe, C.; Jouary, T.; Hauschild, A.; Grob, J.J.; et al. Combined BRAF and MEK Inhibition versus BRAF Inhibition Alone in Melanoma. *N. Engl. J. Med.* **2014**, *371*, 1877–1888. [[CrossRef](#)]
  33. Long, G.V.; Flaherty, K.T.; Stroyakovskiy, D.; Gogas, H.; Levchenko, E.; de Braud, F.; Larkin, J.; Garbe, C.; Jouary, T.; Hauschild, A.; et al. Dabrafenib plus Trametinib versus Dabrafenib Monotherapy in Patients with Metastatic BRAF V600E/K-Mutant Melanoma: Long-Term Survival and Safety Analysis of a Phase 3 Study. *Ann. Oncol.* **2017**, *28*, 1631–1639. [[CrossRef](#)]
  34. Robert, C.; Grob, J.J.; Stroyakovskiy, D.; Karaszewska, B.; Hauschild, A.; Levchenko, E.; Chiarion Sileni, V.; Schachter, J.; Garbe, C.; Bondarenko, I.; et al. Five-Year Outcomes with Dabrafenib plus Trametinib in Metastatic Melanoma. *N. Engl. J. Med.* **2019**, *381*, 626–636. [[CrossRef](#)]
  35. Dummer, R.; Ascierto, P.A.; Gogas, H.J.; Arance, A.; Mandalà, M.; Liskay, G.; Garbe, C.; Schadendorf, D.; Krajsova, I.; Gutzmer, R.; et al. Encorafenib plus Binimetinib versus Vemurafenib or Encorafenib in Patients with BRAF-Mutant Melanoma (COLUMBUS): A Multicentre, Open-Label, Randomised Phase 3 Trial. *Lancet Oncol.* **2018**, *19*, 603–615. [[CrossRef](#)]
  36. Dummer, R.; Ascierto, P.A.; Gogas, H.J.; Arance, A.; Mandalà, M.; Liskay, G.; Garbe, C.; Schadendorf, D.; Krajsova, I.; Gutzmer, R.; et al. Overall Survival in Patients with BRAF-Mutant Melanoma Receiving Encorafenib plus Binimetinib versus Vemurafenib or Encorafenib (COLUMBUS): A Multicentre, Open-Label, Randomised, Phase 3 Trial. *Lancet Oncol.* **2018**, *19*, 1315–1327. [[CrossRef](#)]



37. Dummer, R.; Flaherty, K.T.; Robert, C.; Arance, A.; de Groot, J.W.B.; Garbe, C.; Gogas, H.J.; Gutzmer, R.; Krajsová, I.; Liszkay, G.; et al. COLUMBUS 5-Year Update: A Randomized, Open-Label, Phase III Trial of Encorafenib plus Binimetinib versus Vemurafenib or Encorafenib in Patients With BRAF V600–Mutant Melanoma. *J. Clin. Oncol.* **2022**, *40*, 4178–4188. [\[CrossRef\]](#)
38. Schadendorf, D.; Long, G.V.; Stroiakovski, D.; Karaszewska, B.; Hauschild, A.; Levchenko, E.; Chiarion-Sileni, V.; Schachter, J.; Garbe, C.; Dutriaux, C.; et al. Three-Year Pooled Analysis of Factors Associated with Clinical Outcomes across Dabrafenib and Trametinib Combination Therapy Phase 3 Randomised Trials. *Eur. J. Cancer* **2017**, *82*, 45–55. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Hamid, O.; Cowey, C.L.; Offner, M.; Faries, M.; Carvajal, R.D. Efficacy, Safety, and Tolerability of Approved Combination BRAF and MEK Inhibitor Regimens for BRAF-Mutant Melanoma. *Cancers* **2019**, *11*, 1642. [\[CrossRef\]](#)
40. Grob, J.J.; Amonkar, M.M.; Karaszewska, B.; Schachter, J.; Dummer, R.; Mackiewicz, A.; Stroyakovskiy, D.; Drucis, K.; Grange, F.; Chiarion-Sileni, V.; et al. Comparison of Dabrafenib and Trametinib Combination Therapy with Vemurafenib Monotherapy on Health-Related Quality of Life in Patients with Unresectable or Metastatic Cutaneous BRAF Val600-Mutation-Positive Melanoma (COMBI-v): Results of a Phase 3, Open-Label, Randomised Trial. *Lancet Oncol.* **2015**, *16*, 1389–1398. [\[CrossRef\]](#) [\[PubMed\]](#)
41. Schadendorf, D.; Amonkar, M.M.; Stroyakovskiy, D.; Levchenko, E.; Gogas, H.; de Braud, F.; Grob, J.-J.; Bondarenko, I.; Garbe, C.; Lebbe, C.; et al. Health-Related Quality of Life Impact in a Randomised Phase III Study of the Combination of Dabrafenib and Trametinib versus Dabrafenib Monotherapy in Patients with BRAF V600 Metastatic Melanoma. *Eur. J. Cancer* **2015**, *51*, 833–840. [\[CrossRef\]](#) [\[PubMed\]](#)
42. Lai-Kwon, J.; Inderjeeth, A.J.; Lisy, K.; Sandhu, S.; Rutherford, C.; Jefford, M. Impact of Immune Checkpoint Inhibitors and Targeted Therapy on Health-Related Quality of Life of People with Stage III and IV Melanoma: A Mixed-Methods Systematic Review. *Eur. J. Cancer* **2023**, *184*, 83–105. [\[CrossRef\]](#)
43. Florent, L.; Saby, C.; Slimano, F.; Morjani, H. BRAF V600-Mutated Metastatic Melanoma and Targeted Therapy Resistance: An Update of the Current Knowledge. *Cancers* **2023**, *15*, 2607. [\[CrossRef\]](#) [\[PubMed\]](#)
44. Ng, G.; Xu, W.; Atkinson, V. Treatment Approaches for Melanomas That Relapse After Adjuvant or Neoadjuvant Therapy. *Curr. Oncol. Rep.* **2022**, *24*, 1273–1280. [\[CrossRef\]](#) [\[PubMed\]](#)
45. Ugurel, S.; Röhm, J.; Ascierto, P.A.; Becker, J.C.; Flaherty, K.T.; Grob, J.J.; Hauschild, A.; Larkin, J.; Livingstone, E.; Long, G.V.; et al. Survival of Patients with Advanced Metastatic Melanoma: The Impact of MAP Kinase Pathway Inhibition and Immune Checkpoint Inhibition—Update 2019. *Eur. J. Cancer* **2020**, *130*, 126–138. [\[CrossRef\]](#) [\[PubMed\]](#)
46. Ugurel, S.; Röhm, J.; Ascierto, P.A.; Flaherty, K.T.; Grob, J.J.; Hauschild, A.; Larkin, J.; Long, G.V.; Lorigan, P.; McArthur, G.A.; et al. Survival of Patients with Advanced Metastatic Melanoma: The Impact of Novel Therapies—Update 2017. *Eur. J. Cancer* **2017**, *83*, 247–257. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Thornton, J.; Chhabra, G.; Singh, C.K.; Guzmán-Pérez, G.; Shirley, C.A.; Ahmad, N. Mechanisms of Immunotherapy Resistance in Cutaneous Melanoma: Recognizing a Shapeshifter. *Front Oncol.* **2022**, *12*, 880876. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Pham, J.P.; Joshua, A.M.; da Silva, I.P.; Dummer, R.; Goldinger, S.M. Chemotherapy in Cutaneous Melanoma: Is There Still a Role? *Curr. Oncol. Rep.* **2023**, *25*, 609–621. [\[CrossRef\]](#)
49. Seghers, A.C.; Wilgenhof, S.; Lebbé, C.; Neyns, B. Successful Rechallenge in Two Patients with BRAF-V600-Mutant Melanoma Who Experienced Previous Progression during Treatment with a Selective BRAF Inhibitor. *Melanoma Res.* **2012**, *22*, 466–472. [\[CrossRef\]](#)
50. Romano, E.; Pradervand, S.; Paillusson, A.; Weber, J.; Harshman, K.; Muehlethaler, K.; Speiser, D.; Peters, S.; Rimoldi, D.; Michielin, O. Identification of Multiple Mechanisms of Resistance to Vemurafenib in a Patient with BRAFV600E-Mutated Cutaneous Melanoma Successfully Rechallenged after Progression. *Clin. Cancer Res.* **2013**, *19*, 5749–5757. [\[CrossRef\]](#)
51. Mackiewicz-Wysocka, M.; Krokowicz, Ł.; Kocur, J.; Mackiewicz, J. Resistance to Vemurafenib Can Be Reversible after Treatment Interruption. *Medicine* **2014**, *93*, e157. [\[CrossRef\]](#)
52. Amann, V.C.; Hoffmann, D.; Mangana, J.; Dummer, R.; Goldinger, S.M. Successful Retreatment with Combined BRAF/MEK Inhibition in Metastatic BRAFV600-Mutated Melanoma. *J. Eur. Acad. Dermatol. Venereol.* **2017**, *31*, 1638–1640. [\[CrossRef\]](#) [\[PubMed\]](#)
53. Reschke, R.; Simon, J.; Ziemer, M. Rechallenge of Targeted Therapy in Metastatic Melanoma. *JDDG J. Dtsch. Dermatol. Ges.* **2019**, *17*, 483–486. [\[CrossRef\]](#)
54. Viñal, D.; Martinez, D.; Espinosa, E. Efficacy of Rechallenge with BRAF Inhibition Therapy in Patients with Advanced BRAFV600 Mutant Melanoma. *Clin. Transl. Oncol.* **2019**, *21*, 1061–1066. [\[CrossRef\]](#) [\[PubMed\]](#)
55. Sterne, J.A.; Hernán, M.A.; Reeves, B.C.; Savović, J.; Berkman, N.D.; Viswanathan, M.; Henry, D.; Altman, D.G.; Ansari, M.T.; Boutron, I.; et al. ROBINS-I: A Tool for Assessing Risk of Bias in Non-Randomised Studies of Interventions. *BMJ* **2016**, *355*, i4919. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Wells, G.A.; Shea, B.; O’Connell, D.; Peterson, J.; Welch, V.; Losos, M.; Tugwell, P. The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomized Studies in Meta-Analyses. Available online: [https://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](https://www.ohri.ca/programs/clinical_epidemiology/oxford.asp) (accessed on 16 June 2023).
57. Page, M.J.; McKenzie, J.E.; Bossuyt, P.M.; Boutron, I.; Hoffmann, T.C.; Mulrow, C.D.; Shamseer, L.; Tetzlaff, J.M.; Akl, E.A.; Brennan, S.E.; et al. The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews. *BMJ* **2021**, *372*, n71. [\[CrossRef\]](#) [\[PubMed\]](#)
58. Barker, T.H.; Migliavaca, C.B.; Stein, C.; Colpani, V.; Falavigna, M.; Aromataris, E.; Munn, Z. Conducting Proportional Meta-Analysis in Different Types of Systematic Reviews: A Guide for Synthesisers of Evidence. *BMC Med. Res. Methodol.* **2021**, *21*, 189. [\[CrossRef\]](#)



59. McGrath, S.; Sohn, H.; Steele, R.; Benedetti, A. Meta-analysis of the Difference of Medians. *Biom. J.* **2020**, *62*, 69–98. [\[CrossRef\]](#)
60. Atkinson, V.; Batty, K.; Long, G.V.; Carlino, M.S.; Peters, G.D.; Bhawe, P.; Moore, M.A.; Xu, W.; Brown, L.J.; Arneil, M.; et al. Activity and Safety of Third-Line BRAF-Targeted Therapy (TT) Following First-Line TT and Second-Line Immunotherapy (IT) in Advanced Melanoma. *J. Clin. Oncol.* **2020**, *38*, 10049. [\[CrossRef\]](#)
61. Cybulska-Stopa, B.; Rogala, P.; Czarnecka, A.M.; Galus, Ł.; Dziura, R.; Rajczykowski, M.; Kubiowski, T.; Wiśniewska, M.; Gega-Czarnota, A.; Teterycz, P.; et al. BRAF and MEK Inhibitors Rechallenge as Effective Treatment for Patients with Metastatic Melanoma. *Melanoma Res.* **2020**, *30*, 465–471. [\[CrossRef\]](#)
62. Persa, O.D.; Mauch, C. Outcomes after Retreatment with MAPK Inhibitors and Immune Checkpoint Inhibitors in Melanoma Patients. *Future Oncol.* **2021**, *17*, 3809–3817. [\[CrossRef\]](#)
63. Roux, J.; Pages, C.; Malouf, D.; Basset Seguin, N.; Madjlessi, N.; Baccard, M.; Comte, C.; Archimbaud, A.; Battistella, M.; Viguier, M.; et al. BRAF Inhibitor Rechallenge in Patients with Advanced BRAF V600-Mutant Melanoma. *Melanoma Res.* **2015**, *25*, 559–563. [\[CrossRef\]](#)
64. Schreuer, M.; Jansen, Y.; Planken, S.; Chevolet, I.; Seremet, T.; Kruse, V.; Neyns, B. Combination of Dabrafenib plus Trametinib for BRAF and MEK Inhibitor Pretreated Patients with Advanced BRAFV600-Mutant Melanoma: An Open-Label, Single Arm, Dual-Centre, Phase 2 Clinical Trial. *Lancet Oncol.* **2017**, *18*, 464–472. [\[CrossRef\]](#)
65. Tietze, J.K.; Forschner, A.; Loquai, C.; Mitzel-Rink, H.; Zimmer, L.; Meiss, F.; Rafei-Shamsabadi, D.; Utikal, J.; Bergmann, M.; Meier, F.; et al. The Efficacy of Re-Challenge with BRAF Inhibitors after Previous Progression to BRAF Inhibitors in Melanoma: A Retrospective Multicenter Study. *Oncotarget* **2018**, *9*, 34336–34346. [\[CrossRef\]](#)
66. Valpione, S.; Carlino, M.S.; Mangana, J.; Mooradian, M.J.; McArthur, G.; Schadendorf, D.; Hauschild, A.; Menzies, A.M.; Arance, A.; Ascierto, P.A.; et al. Rechallenge with BRAF-Directed Treatment in Metastatic Melanoma: A Multi-Institutional Retrospective Study. *Eur. J. Cancer* **2018**, *91*, 116–124. [\[CrossRef\]](#)
67. Chapman, P.B.; Robert, C.; Larkin, J.; Haanen, J.B.; Ribas, A.; Hogg, D.; Hamid, O.; Ascierto, P.A.; Testori, A.; Lorigan, P.C.; et al. Vemurafenib in Patients with BRAFV600 Mutation-Positive Metastatic Melanoma: Final Overall Survival Results of the Randomized BRIM-3 Study. *Ann. Oncol.* **2017**, *28*, 2581–2587. [\[CrossRef\]](#)
68. Hauschild, A.; Ascierto, P.A.; Schadendorf, D.; Grob, J.J.; Ribas, A.; Kiecker, F.; Dutriaux, C.; Demidov, L.V.; Lebbé, C.; Rutkowski, P.; et al. Long-Term Outcomes in Patients with BRAF V600-Mutant Metastatic Melanoma Receiving Dabrafenib Monotherapy: Analysis from Phase 2 and 3 Clinical Trials. *Eur. J. Cancer* **2020**, *125*, 114–120. [\[CrossRef\]](#)
69. Long, G.V.; Stroyakovskiy, D.; Gogas, H.; Levchenko, E.; de Braud, F.; Larkin, J.; Garbe, C.; Jouary, T.; Hauschild, A.; Grob, J.-J.; et al. Dabrafenib and Trametinib versus Dabrafenib and Placebo for Val600 BRAF-Mutant Melanoma: A Multicentre, Double-Blind, Phase 3 Randomised Controlled Trial. *Lancet* **2015**, *386*, 444–451. [\[CrossRef\]](#) [\[PubMed\]](#)
70. Savoia, P.; Zavattaro, E.; Cremona, O. Clinical Implications of Acquired BRAF Inhibitors Resistance in Melanoma. *Int. J. Mol. Sci.* **2020**, *21*, 9730. [\[CrossRef\]](#) [\[PubMed\]](#)
71. Ascierto, P.A.; Dummer, R.; Gogas, H.J.; Flaherty, K.T.; Arance, A.; Mandala, M.; Liskay, G.; Garbe, C.; Schadendorf, D.; Krajsova, I.; et al. Update on Tolerability and Overall Survival in COLUMBUS: Landmark Analysis of a Randomised Phase 3 Trial of Encorafenib plus Binimetinib vs. Vemurafenib or Encorafenib in Patients with BRAF V600-Mutant Melanoma. *Eur. J. Cancer* **2020**, *126*, 33–44. [\[CrossRef\]](#)
72. Tian, Y.; Guo, W. A Review of the Molecular Pathways Involved in Resistance to BRAF Inhibitors in Patients with Advanced-Stage Melanoma. *Med. Sci. Monit.* **2020**, *26*, e920957-1. [\[CrossRef\]](#)
73. Nazarian, R.; Shi, H.; Wang, Q.; Kong, X.; Koya, R.C.; Lee, H.; Chen, Z.; Lee, M.-K.; Attar, N.; Sazegar, H.; et al. Melanomas Acquire Resistance to B-RAF(V600E) Inhibition by RTK or N-RAS Upregulation. *Nature* **2010**, *468*, 973–977. [\[CrossRef\]](#)
74. Johnson, D.B.; Menzies, A.M.; Zimmer, L.; Eroglu, Z.; Ye, F.; Zhao, S.; Rizos, H.; Sucker, A.; Scolyer, R.A.; Gutzmer, R.; et al. Acquired BRAF Inhibitor Resistance: A Multicenter Meta-Analysis of the Spectrum and Frequencies, Clinical Behaviour, and Phenotypic Associations of Resistance Mechanisms. *Eur. J. Cancer* **2015**, *51*, 2792–2799. [\[CrossRef\]](#)
75. Sun, C.; Wang, L.; Huang, S.; Heynen, G.J.J.E.; Prahallad, A.; Robert, C.; Haanen, J.; Blank, C.; Wesseling, J.; Willems, S.M.; et al. Reversible and Adaptive Resistance to BRAF(V600E) Inhibition in Melanoma. *Nature* **2014**, *508*, 118–122. [\[CrossRef\]](#)
76. Slominski, R.M.; Raman, C.; Chen, J.Y.; Slominski, A.T. How Cancer Hijacks the Body's Homeostasis through the Neuroendocrine System. *Trends Neurosci.* **2023**, *46*, 263–275. [\[CrossRef\]](#) [\[PubMed\]](#)
77. Slominski, R.M.; Sarna, T.; Płonka, P.M.; Raman, C.; Brożyna, A.A.; Slominski, A.T. Melanoma, Melanin, and Melanogenesis: The Yin and Yang Relationship. *Front Oncol.* **2022**, *12*, 842496. [\[CrossRef\]](#)
78. Smith, L.K.; Parmenter, T.; Kleinschmidt, M.; Kusnadi, E.P.; Kang, J.; Martin, C.A.; Lau, P.; Patel, R.; Lorent, J.; Papadopoulos, D.; et al. Adaptive Translational Reprogramming of Metabolism Limits the Response to Targeted Therapy in BRAFV600 Melanoma. *Nat. Commun.* **2022**, *13*, 1100. [\[CrossRef\]](#)
79. Yan, C.; Chen, S.C.; Ayers, G.D.; Nebhan, C.A.; Roland, J.T.; Weiss, V.L.; Johnson, D.B.; Richmond, A. Proximity of Immune and Tumor Cells Underlies Response to BRAF/MEK-Targeted Therapies in Metastatic Melanoma Patients. *NPJ Precis. Oncol.* **2022**, *6*, 6. [\[CrossRef\]](#) [\[PubMed\]](#)
80. Zuo, Q.; Liu, J.; Huang, L.; Qin, Y.; Hawley, T.; Seo, C.; Merlino, G.; Yu, Y. AXL/AKT Axis Mediated-Resistance to BRAF Inhibitor Depends on PTEN Status in Melanoma. *Oncogene* **2018**, *37*, 3275–3289. [\[CrossRef\]](#) [\[PubMed\]](#)

81. McGill, G.G.; Horstmann, M.; Widlund, H.R.; Du, J.; Motyckova, G.; Nishimura, E.K.; Lin, Y.-L.; Ramaswamy, S.; Avery, W.; Ding, H.-F.; et al. Bcl2 Regulation by the Melanocyte Master Regulator Mitf Modulates Lineage Survival and Melanoma Cell Viability. *Cell* **2002**, *109*, 707–718. [\[CrossRef\]](#) [\[PubMed\]](#)
82. Konieczkowski, D.J.; Johannessen, C.M.; Abudayyeh, O.; Kim, J.W.; Cooper, Z.A.; Piris, A.; Frederick, D.T.; Barzily-Rokni, M.; Straussman, R.; Haq, R.; et al. A Melanoma Cell State Distinction Influences Sensitivity to MAPK Pathway Inhibitors. *Cancer Discov.* **2014**, *4*, 816–827. [\[CrossRef\]](#)
83. Khaliq, M.; Manikkam, M.; Martinez, E.D.; Fallahi-Sichani, M. Epigenetic Modulation Reveals Differentiation State Specificity of Oncogene Addiction. *Nat. Commun.* **2021**, *12*, 1536. [\[CrossRef\]](#) [\[PubMed\]](#)
84. Karras, P.; Bordeu, I.; Pozniak, J.; Nowosad, A.; Pazzi, C.; Van Raemdonck, N.; Landeloos, E.; Van Herck, Y.; Pedri, D.; Bervoets, G.; et al. A Cellular Hierarchy in Melanoma Uncouples Growth and Metastasis. *Nature* **2022**, *610*, 190–198. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Centeno, P.P.; Pavet, V.; Marais, R. The Journey from Melanocytes to Melanoma. *Nat. Rev. Cancer* **2023**, *23*, 372–390. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Frederick, D.T.; Piris, A.; Cogdill, A.P.; Cooper, Z.A.; Lezcano, C.; Ferrone, C.R.; Mitra, D.; Boni, A.; Newton, L.P.; Liu, C.; et al. BRAF Inhibition Is Associated with Enhanced Melanoma Antigen Expression and a More Favorable Tumor Microenvironment in Patients with Metastatic Melanoma. *Clin. Cancer Res.* **2013**, *19*, 1225–1231. [\[CrossRef\]](#)
87. Wilmott, J.S.; Long, G.V.; Howle, J.R.; Haydu, L.E.; Sharma, R.N.; Thompson, J.F.; Kefford, R.F.; Hersey, P.; Scolyer, R.A. Cancer Therapy: Clinical Selective BRAF Inhibitors Induce Marked T-Cell Infiltration into Human Metastatic Melanoma. *Clin. Cancer Res.* **2012**, *18*, 1386–1394. [\[CrossRef\]](#) [\[PubMed\]](#)
88. Stege, H.; Haist, M.; Nikfarjam, U.; Schultheis, M.; Heinz, J.; Pémmler, S.; Loquai, C.; Grabbe, S. The Status of Adjuvant and Neoadjuvant Melanoma Therapy, New Developments and Upcoming Challenges. *Target. Oncol.* **2021**, *16*, 537–552. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Mason, R.; Dearden, H.C.; Nguyen, B.; Soon, J.A.; Smith, J.L.; Randhawa, M.; Mant, A.; Warburton, L.; Lo, S.; Meniawy, T.; et al. Combined Ipilimumab and Nivolumab First-line and after BRAF-targeted Therapy in Advanced Melanoma. *Pigment. Cell Melanoma Res.* **2020**, *33*, 358–365. [\[CrossRef\]](#)
90. Haas, L.; Elewaut, A.; Gerard, C.L.; Umkehrer, C.; Leiendecker, L.; Pedersen, M.; Krecioch, I.; Hoffmann, D.; Novatchkova, M.; Kuttke, M.; et al. Acquired Resistance to Anti-MAPK Targeted Therapy Confers an Immune-Evasive Tumor Microenvironment and Cross-Resistance to Immunotherapy in Melanoma. *Nat. Cancer* **2021**, *2*, 693–708. [\[CrossRef\]](#)
91. Wang, Y.; Liu, S.; Yang, Z.; Algazi, A.P.; Lomeli, S.H.; Wang, Y.; Othus, M.; Hong, A.; Wang, X.; Randolph, C.E.; et al. Anti-PD-1/L1 Lead-in before MAPK Inhibitor Combination Maximizes Antitumor Immunity and Efficacy. *Cancer Cell* **2021**, *39*, 1375–1387.e6. [\[CrossRef\]](#)
92. Chandra, S.; Choi, J.S.; Sosman, J.A. Melanoma: Does Sequencing Really Matter? *J. Clin. Oncol.* **2023**, *41*, 167–169. [\[CrossRef\]](#)
93. Ascierto, P.A.; Mandalà, M.; Ferrucci, P.F.; Guidoboni, M.; Rutkowski, P.; Ferraresi, V.; Arance, A.; Guida, M.; Maiello, E.; Gogas, H.; et al. Sequencing of Ipilimumab Plus Nivolumab and Encorafenib plus Binimetinib for Untreated BRAF-Mutated Metastatic Melanoma (SECOMBIT): A Randomized, Three-Arm, Open-Label Phase II Trial. *J. Clin. Oncol.* **2023**, *41*, 212–221. [\[CrossRef\]](#)
94. Atkins, M.B.; Lee, S.J.; Chmielowski, B.; Tarhini, A.A.; Cohen, G.I.; Truong, T.G.; Moon, H.H.; Davar, D.; O'Rourke, M.; Stephenson, J.J.; et al. Combination Dabrafenib and Trametinib Versus Combination Nivolumab and Ipilimumab for Patients with Advanced BRAF-Mutant Melanoma: The DREAMseq Trial—ECOG-ACRIN EA6134. *J. Clin. Oncol.* **2023**, *41*, 186–197. [\[CrossRef\]](#)
95. Welte, M.; Dimitriou, F.; Gutzmer, R.; Dummer, R. Triple Combination of Immune Checkpoint Inhibitors and BRAF/MEK Inhibitors in BRAF V600 Melanoma: Current Status and Future Perspectives. *Cancers* **2022**, *14*, 5489. [\[CrossRef\]](#)
96. Switzer, B.; Puzanov, I.; Skitzki, J.J.; Hamad, L.; Ernstoff, M.S. Managing Metastatic Melanoma in 2022: A Clinical Review. *JCO Oncol. Pract.* **2022**, *18*, 335–351. [\[CrossRef\]](#)
97. van Breeschoten, J.; Wouters, M.W.J.M.; Hilarius, D.L.; Haanen, J.B.; Blank, C.U.; Aarts, M.J.B.; van den Bergmortel, F.W.P.J.; de Groot, J.W.B.; Hospers, G.A.P.; Kapiteijn, E.; et al. First-Line BRAF/MEK Inhibitors versus Anti-PD-1 Monotherapy in BRAFV600-Mutant Advanced Melanoma Patients: A Propensity-Matched Survival Analysis. *Br. J. Cancer* **2021**, *124*, 1222–1230. [\[CrossRef\]](#)
98. Adeleke, S.; Okoli, S.; Augustine, A.; Galante, J.; Agnihotri, A.; Uccello, M.; Ghose, A.; Moschetta, M.; Boussios, S. Melanoma—The Therapeutic Considerations in the Clinical Practice. *Ann. Palliat. Med.* **2023**. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Fernandez, M.F.; Choi, J.; Sosman, J. New Approaches to Targeted Therapy in Melanoma. *Cancers* **2023**, *15*, 3224. [\[CrossRef\]](#) [\[PubMed\]](#)
100. Moreno-Ramírez, D.; Fernández-Orland, A.; Ferrándiz, L. Immunotherapy and Targeted Therapy in Patients With Advanced Melanoma and the V600 BRAF Mutation: Which One First? *Actas. Dermosifiliogr.* **2023**. [\[CrossRef\]](#)
101. Das Thakur, M.; Salangsang, F.; Landman, A.S.; Sellers, W.R.; Pryer, N.K.; Levesque, M.P.; Dummer, R.; McMahon, M.; Stuart, D.D. Modelling Vemurafenib Resistance in Melanoma Reveals a Strategy to Forestall Drug Resistance. *Nature* **2013**, *494*, 251–255. [\[CrossRef\]](#)
102. Kuczyński, E.A.; Sargent, D.J.; Grothey, A.; Kerbel, R.S. Drug Rechallenge and Treatment beyond Progression—Implications for Drug Resistance. *Nat. Rev. Clin. Oncol.* **2013**, *10*, 571–587. [\[CrossRef\]](#)
103. Gebhardt, C.; Ascierto, P.; Atkinson, V.; Corrie, P.; Dummer, R.; Schadendorf, D. The Concepts of Rechallenge and Retreatment in Melanoma: A Proposal for Consensus Definitions. *Eur. J. Cancer* **2020**, *138*, 68–76. [\[CrossRef\]](#)

104. Devji, T.; Levine, O.; Neupane, B.; Beyene, J.; Xie, F. Systemic Therapy for Previously Untreated Advanced BRAF-Mutated Melanoma. *JAMA Oncol.* **2017**, *3*, 366. [[CrossRef](#)] [[PubMed](#)]
105. Seth, R.; Messersmith, H.; Kaur, V.; Kirkwood, J.M.; Kudchadkar, R.; McQuade, J.L.; Provenzano, A.; Swami, U.; Weber, J.; Alluri, K.C.; et al. Systemic Therapy for Melanoma: ASCO Guideline. *J. Clin. Oncol.* **2020**, *38*, 3947–3970. [[CrossRef](#)] [[PubMed](#)]
106. Michielin, O.; van Akkooi, A.C.J.; Ascierto, P.A.; Dummer, R.; Keilholz, U. Cutaneous Melanoma: ESMO Clinical Practice Guidelines for Diagnosis, Treatment and Follow-Up. *Ann. Oncol.* **2019**, *30*, 1884–1901. [[CrossRef](#)] [[PubMed](#)]
107. Fujisawa, Y.; Ito, T.; Kato, H.; Irie, H.; Kaji, T.; Maekawa, T.; Asai, J.; Yamamoto, Y.; Fujimura, T.; Nakai, Y.; et al. Outcome of Combination Therapy Using BRAF and MEK Inhibitors among Asian Patients with Advanced Melanoma: An Analysis of 112 Cases. *Eur. J. Cancer* **2021**, *145*, 210–220. [[CrossRef](#)] [[PubMed](#)]
108. Colombino, M.; Capone, M.; Lissia, A.; Cossu, A.; Rubino, C.; De Giorgi, V.; Massi, D.; Fonsatti, E.; Staibano, S.; Nappi, O.; et al. BRAF/NRAS Mutation Frequencies Among Primary Tumors and Metastases in Patients With Melanoma. *J. Clin. Oncol.* **2012**, *30*, 2522–2529. [[CrossRef](#)]
109. Soffietti, R.; Ahluwalia, M.; Lin, N.; Rudà, R. Management of Brain Metastases According to Molecular Subtypes. *Nat. Rev. Neurol.* **2020**, *16*, 557–574. [[CrossRef](#)] [[PubMed](#)]
110. Nakamura, Y.; Ishitsuka, Y.; Tanaka, R.; Okiyama, N.; Watanabe, R.; Saito, A.; Furuta, J.; Fujisawa, Y. Frequent Brain Metastases during Treatment with BRAF/MEK Inhibitors: A Retrospective Single Institutional Study. *J. Dermatol.* **2020**, *47*, 1191–1194. [[CrossRef](#)]
111. Wagner, N.B.; Forschner, A.; Leiter, U.; Garbe, C.; Eigentler, T.K. S100B and LDH as Early Prognostic Markers for Response and Overall Survival in Melanoma Patients Treated with Anti-PD-1 or Combined Anti-PD-1 plus Anti-CTLA-4 Antibodies. *Br. J. Cancer* **2018**, *119*, 339–346. [[CrossRef](#)] [[PubMed](#)]
112. Internò, V.; Sergi, M.C.; Metta, M.E.; Guida, M.; Trerotoli, P.; Strippoli, S.; Circelli, S.; Porta, C.; Tucci, M. Melanoma Brain Metastases: A Retrospective Analysis of Prognostic Factors and Efficacy of Multimodal Therapies. *Cancers* **2023**, *15*, 1542. [[CrossRef](#)]
113. Ascierto, P.A.; Ribas, A.; Larkin, J.; McArthur, G.A.; Lewis, K.D.; Hauschild, A.; Flaherty, K.T.; McKenna, E.; Zhu, Q.; Mun, Y.; et al. Impact of Initial Treatment and Prognostic Factors on Postprogression Survival in BRAF-Mutated Metastatic Melanoma Treated with Dacarbazine or Vemurafenib ± Cobimetinib: A Pooled Analysis of Four Clinical Trials. *J. Transl. Med.* **2020**, *18*, 294. [[CrossRef](#)]
114. Heinzerling, L.; Eigentler, T.K.; Fluck, M.; Hassel, J.C.; Heller-Schenck, D.; Leipe, J.; Pauschinger, M.; Vogel, A.; Zimmer, L.; Gutzmer, R. Tolerability of BRAF/MEK Inhibitor Combinations: Adverse Event Evaluation and Management. *ESMO Open* **2019**, *4*, e000491. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.